

# On nonlocal modified gravity with cosmological solutions

Ivan Dimitrijevic

Faculty of Mathematics, University of Belgrade  
Studentski trg 16, Belgrade, Serbia

In this talk, we consider nonlocal gravity action without matter in the form

$$S = \int d^4x \sqrt{-g} \left( \frac{R - 2\Lambda}{16\pi G} + R^p \mathcal{F}(\square) R \right),$$

where  $\mathcal{F}(\square)$  is an analytic function of the d'Alembertian  $\square$  and  $p = +1, -1$ . We present a few  $a(t)$  nonsingular bounce cosmological solutions for the above two actions using  $FLRW$  metric. see references [1-5].

[1] T. Biswas, T. Koivisto, A. Mazumdar, “Towards a resolution of the cosmological singularity in non-local higher derivative theories of gravity”, JCAP **1011** (2010) 008 [arXiv:1005.0590v2 [hep-th]]. [2] A. S. Koshelev, S. Yu. Vernov, “On bouncing solutions in non-local gravity”, Phys. Part. Nuclei **43**, 666–668 (2012)[arXiv:1202.1289v1 [hep-th]]. [3] T. Biswas, A. S. Koshelev, A. Mazumdar, S. Yu. Vernov, “Stable bounce and inflation in non-local higher derivative cosmology”, JCAP **08** (2012) 024, [arXiv:1206.6374v2 [astro-ph.CO]]. [4] I. Dimitrijevic, B. Dragovich, J. Grujic, Z. Rakic, “New cosmological solutions in nonlocal modified gravity”, Rom. Journ. Phys. **58** (5-6), 550–559 (2013) [arXiv:1302.2794 [gr-qc]]. [5] I. Dimitrijevic, B. Dragovich, J. Grujic, Z. Rakic, “A new model of nonlocal modified gravity”, Publications de l’Institut Mathematique **94** (108), 187–196 (2013).

This is joint work with B. Dragovich, J. Grujic and Z. Rakic.